

ORIGINAL PAPER



WILEY

Parents who inject drugs: Demographics, care arrangements and correlates for child placement in out-of-home care

Jocelyn Chan¹ | Bernadette Ward^{1,2} | Lisa Maher^{1,3} | Sione Crawford^{1,4} | Mark Stoové^{1,5} | Paul Dietze^{1,6}

¹Burnet Institute, Melbourne, Australia

²School of Rural Health, Monash University, Melbourne, Australia

³Kirby Institute, UNSW Sydney, Sydney, Australia

⁴Harm Reduction Victoria, Melbourne, Australia

⁵School of Public Health and Preventive Medicine, Monash University, Melbourne, Australia

⁶National Drug Research Institute, Curtin University, Melbourne, Australia

Correspondence

Jocelyn Chan, Burnet Institute,
85 Commercial Road, Melbourne, VIC
3004, Australia.
Email: jocelyn.chan@burnet.edu.au

Funding information

Colonial Foundation; National Health and Medical Research Council, Grant/Award Numbers: 1126090, 545891; Gilead Sciences; AbbVie; Victorian Operational Infrastructure Fund

Abstract

Introduction: Children in families affected by substance use disorders are at high risk of being placed in out-of-home care (OOHC). We aimed to describe the characteristics of parents who inject drugs and identify correlates associated with child placement in OOHC.

Methods: We used baseline data from a community-based cohort of parents who inject drugs (SuperMIX) from Melbourne, Australia. Participants were recruited via convenience, respondent-driven and snowball sampling from April 2008 to November 2020, with follow-up until March 2021. To explore correlates associated with child placement to OOHC, we used multivariable logistic regression and assessed for potential interactions between gender and a range of relevant covariates.

Results: Of the 1067 participants, 611 (57%) reported being parents. Fifty-six percent of parents reported child protection involvement. Almost half (49%) had children in OOHC. Nearly half of the parents lived in unstable accommodation (44%) and many of them experienced moderate–severe levels of anxiety (48%) and depression (53%). Female or non-binary gender, identifying as Aboriginal or Torres Strait Islander, experiencing assault and having more children were associated with child removal to OOHC. Of the 563 participants who reported their own childhood care status, 135 (24%) reported they had been removed to OOHC.

Discussion and Conclusions: We identified high rates of child placement in OOHC among parents who inject drugs. There is a need for targeted health and social services, that are gender and culturally responsive, in addition to systems-level interventions addressing social inequities, such as housing, to support parents to care for their children.

KEYWORDS

child protection, injecting drug use, out-of-home care, parenting

Key points

- We identified high rates of child protection involvement and child removal to out-of-home-care (OOHC) among a cohort of parents of inject drugs.

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2023 The Authors. *Drug and Alcohol Review* published by John Wiley & Sons Australia, Ltd on behalf of Australasian Professional Society on Alcohol and other Drugs.

- Parents who themselves had been removed to OOHC as a child were more likely to have child protection involvement and OOHC arrangements with their own children.
- Parents who injected drugs had high psychosocial needs, including high rates of moderate–severe anxiety and depression and unstable housing.
- Being female or non-binary, identifying as Aboriginal or Torres Strait Islander, experiencing assault and having a greater number of children were associated with child removal to OOHC.
- Further research and intervention are urgently needed to support parents to care for their children, given the intergenerational impacts of child removal to OOHC.

1 | INTRODUCTION

Globally, an estimated 15.6 million people aged 15–64 years inject drugs in 2016, many of whom are parents [1]. With sustained use of alcohol and other drugs, substance use disorders, characterised by loss of control over the use and subsequent impact on a person's life, can develop. In 2016, the global prevalence of opioid and amphetamine use disorders was estimated at 353 and 65 per 100,000 people, respectively [2].

Children in families affected by substance use disorders are at high risk of being placed in out-of-home care (OOHC) [3]. It is estimated that 60% of Australian children in OOHC were removed from households affected by substance use disorders [4]. For example, among a cohort of 171 Australian mothers in treatment for opioid use disorder, 64% had been reported to child protection services and 33% had at least one child in OOHC [5]. While, differing definitions of OOHC by jurisdiction and over time care limits comparability, data indicate rates of OOHC have been rising in Australia over the last decade. In the past 5 years, overall rates of OOHC have stabilised, but continue to increase for Aboriginal and Torres Strait Islander families who remain over-represented alongside families affected by substance use disorders [6, 7].

Preventing entry to OOHC or mitigating its impacts is critical given the long-term implications for both parents and children. Systematic reviews of non-experimental studies have consistently found poorer health and social outcomes for children placed in OOHC compared to their peers [8, 9]. Evidence indicates that patterns of OOHC can continue across generations, mediated through mental health problems, particularly among families experiencing socioeconomic disadvantage [10]. However, there is evidence that protective factors, including supportive relationships and educational attainment can disrupt the intergenerational transmission of placement in OOHC [11, 12]. Evidence also indicates that loss of child custody also negatively affects maternal health—with

increased rates of substance use, overdose, mental illness, trauma and mortality [13, 14].

While children of people who use drugs are often considered to be at risk of poor child outcomes, prior research cautions against conflating parental drug use with child neglect [15–17]. After all, many children with family histories of drug use reach adulthood without demonstrating maladaptive behaviours [18]. Furthermore, any associations between parental drug use and child outcomes need to account for the social contexts and processes which mediate the ability of parents who inject drugs to provide appropriate care to their children. Injection drug use often co-occurs with many other indicators of marginalisation, such as poverty, unstable housing, involvement with the criminal justice system, mental health issues and family conflict, which may impede child development [19, 20]. Additionally, stigmatising attitudes towards people who inject drugs, including parents and especially mothers, create additional barriers to accessing care and social services and increase the likelihood of referral to child protection services and the removal of parental rights [17, 21].

Prior research indicates gendered differences in patterns of child removal among parents who use drugs, with mothers more likely to have children placed in OOHC [22, 23]. This may be due to the higher likelihood of mothers being primary caregivers; however, women also report experiencing amplified stigma because drug use conflicts with normative social expectations for women as caregivers [21, 24]. Gender-specific policies, such as the mandatory reporting of prenatal opioid agonist treatment to child welfare services, reinforce stigma and increase surveillance among women and mothers [25].

Parenthood represents a key time point in which to intervene to improve health outcomes for both parents and children from families affected by the harms associated with substance use. Mothers who use heroin employed a range of strategies to shield their children from the harms of their substance use, including cessation of drug use, entering treatment and ensuring a stable

home environment [16]. Indeed, pregnancy and parenthood have been identified as significant motivators for seeking drug treatment [5]. Conversely, being a parent with custody of a child may be a barrier to accessing treatment given legitimate fears of child removal [22].

Evidence remains scarce on the needs of parents who use illicit substances, including parents who inject drugs. A systematic review identified only two relevant studies describing correlates for child removal among parents who inject drugs [26] and most research to date has been conducted among parents accessing treatment for substance use disorders [27].

Evidence, including the characteristics and living circumstances of families, is needed to develop strategies to support families affected by substance use and facilitate family preservation. The Melbourne Injecting Drug User Cohort Study (SuperMIX) provides a unique opportunity to examine parent characteristics among a community-based cohort of people who inject drugs. The aim of our study was to describe the socio-demographic, health, drug use and parenting characteristics of SuperMIX participants and how these were associated with the removal of their children to OOHC. Given the gendered norms and expectations around parenting roles, we investigated for potential interactions by gender, to assess whether the association between these characteristics and child removal to out-of-home care differed by gender. We also examined the intergenerational association between parents' own childhood OOHC status (i.e. whether they were removed to OOHC) and that of their children.

2 | METHODS

2.1 | Study design, setting and study population

We used data from the SuperMIX study [28]. In summary, participants were recruited through a combination of respondent-driven sampling, snowball sampling and peer outreach from April 2008 to November 2020, with follow-up until March 2021. SuperMIX eligibility criteria included being age over 18 years, injected either heroin or methamphetamine at least six times in the past 6 months, and resident in Melbourne, Victoria. All participants provided written informed consent and were remunerated for study visits. Further details about the cohort and eligibility are reported elsewhere [28]. Ethics approval was received from the Victorian Department of Health Human Research Ethics Committee (approval number: 28.13.17) and the AIHW Ethics Committee (approval number: EO2013/3/48).

2.2 | Data collection

SuperMIX includes interviewer-administered questionnaires covering four main domains: socio-demographic characteristics, drug use characteristics, health and health service utilisation. Parenting questions were introduced in 2012 and asked annually for all participants. In 2017, a question about each participant's own childhood care history was introduced. For this analysis, we used participants' baseline survey responses to the parenting questions and the 2017 responses about their own childhood care experience.

2.3 | Outcome

The primary outcome variable of placement of at least one child currently in OOHC was determined by self-report. We assumed that the participants had understood the question in a way that aligns with the Australian definition of OOHC—that is overnight care for children under the age of 18 years who are deemed unable to live with their family because of safety concerns [6].

Participants were asked whether they had ever been a parent or caregiver. Other self-reported parenting data collected included the number of children, type of relationship, each child's co-residential status and whether child protection had ever been involved in their child(ren)'s care.

2.4 | Covariates

Socio-demographic characteristics examined included age, gender, Aboriginal and/or Torres Strait Islander status, employment status, accommodation status, level of education, co-resident spouse or partner, experience of assault and incarceration history. Drug use characteristics included age at first injection, main illicit drug of choice, frequency of injection and alcohol use (Alcohol Use Disorder Identification Test-C).

The Patient Health Questionnaire and the Generalised Anxiety Disorder (GAD-7) instruments were introduced in 2015 and used to measure depression and anxiety with ranges 0–4 and 5+ categorised as none-mild and moderate-severe depression/anxiety disorder, respectively [29]. Healthcare utilisation was indicated by self-reports of current drug treatment (opioid agonist treatment, OAT) and non-OAT primary care utilisation (general practice) within the past 12 months. Data on participant's own childhood care history included whether they were cared for by at least one parent, whether they were removed to out-of-home care as a

child and whether they were a member or child of the stolen generation. The stolen generation refers to a cohort of Aboriginal and Torres Strait Islander children who were forcibly removed from their families by governments, churches and welfare bodies to be raised in institutions or fostered or adopted by non-Aboriginal or Torres Strait Islander families [30].

A detailed description of key variables used in the analysis is provided in Table S1.

2.5 | Data analysis

We described the socio-demographic, drug use, health-care utilisation and care history characteristics of participants stratified by parent or caregiver status. Differences in characteristics were tested using Pearson's chi square test for categorical variables or the Kruskal–Wallis test for continuous variables. For parents and caregivers, we described the care arrangements of their children stratified by parents' gender. We presented continuous variables with median and interquartile range and categorical variables with number and proportion.

To explore correlates associated with child placement to OOHC, we used bivariable and multivariable logistic regression models. Seventeen covariables were included in the multivariable model (see Table S1). Variables about participant's care history and anxiety/depression scores were excluded as they were not available until 2017 or 2015 onwards, respectively. We assessed potential interactions between gender and a range of relevant covariates (age, Aboriginal and Torres Strait Islander status, employment status, accommodation status, living with spouse or partner, incarceration, experience of assault, current OAT and number of children) using the likelihood ratio test to determine whether the addition of an interaction term yielded a significantly more accurate model [31]. A complete case approach to the regression analyses was adopted where participants with any missing data were excluded. Adjusted odds ratios with 95% confidence intervals were generated with statistical significance set at $p < 0.05$. All analyses were conducted using R Version 4.2.1 statistical software [32]. To assist with the interpretation of interactions, we predicted the odds of removal to OOHC by parents' gender for relevant variables based on the fitted model using the 'avg_predictions' function from the marginal effects package [33].

2.6 | Sub-analysis

A sub-analysis was conducted to determine the association between a parent's own childhood history OOHC and

subsequent placement of their children in OOHC, given these data were only available for a limited subset of participants from 2017 onwards. We replicated the multivariable logistic regression model described above and added parent's own OOHC status as an additional covariate.

3 | RESULTS

Questions about parenting were asked of 1067 unique participants. Among these, the median age was 34 years (interquartile range 23.2–44.8), 358 (34%) were female or non-binary and 142 (13%) participants identified as Aboriginal and/or Torres Strait Islander. For the subset of participants recruited from June 2017 onwards with additional information about the participant's own care histories, 135 out of 563 (24%) reported being removed to out-of-home care as a child with the proportion increasing to 47% ($n = 55$) among Aboriginal and Torres Strait Islander participants. Fifty-four out of 89 Aboriginal and/or Torres Strait Islander participants (61%) reported being either a member or a child of the stolen generation (Table 1).

Of the 1067 participants, 611 (57%) reported being a parent or caregiver (hereafter referred to as parents). The vast majority reported being birth parents to at least one child ($n = 588$, 96%). Parents were more likely to be female, non-binary and Aboriginal or Torres Strait Islander, to have lower educational attainment, be unemployed, currently live in stable accommodation and to have experienced incarceration compared to participants who were not parents (Table 1). Further breakdown of demographic characteristics by both gender and parental status is available in Table S2.

A minority of parents reported residing with at least one of their children ($n = 160$, 26%), although more women and non-binary parents resided with at least one child (38%) compared to men (18%). More than half ($n = 339$, 56%) of parents reported child protection service involvement, increasing to 72% for female or non-binary parents. Almost half ($n = 220$, 49%) of parents who responded reported that their children were currently in OOHC (Table 2). Compared to parents who reported child protection involvement, but did not have their child(ren) removed, parents who had a child removed were more likely to be women, identify as Aboriginal and Torres Strait Islander, report a history of assault and have experienced out-of-home care themselves (Table S3).

3.1 | Correlates of child removal to out-of-home care

Complete case analysis for the multivariable analysis included 391 parents of 965 children. The odds of having

TABLE 1 Characteristics of participants stratified by parent or caregiver status.

	Not parent or caregiver (<i>N</i> = 456)	Parent or caregiver (<i>N</i> = 611)	Overall (<i>N</i> = 1067) ^b	<i>p</i> -value
Gender ^a				<0.001
Male	352 (77.2%)	356 (58.4%)	708 (66.4%)	
Female or non-binary	104 (22.8%)	254 (41.6%)	358 (33.6%)	
Age at interview	33.0 (9.00)	34.0 (12.0)	34.0 (11.0)	<0.001
Aboriginal and Torres Strait Islander				<0.001
Not Aboriginal and/or Torres Strait Islander	425 (93.2%)	497 (81.7%)	922 (86.7%)	
Aboriginal or Torres Strait Islander	31 (6.80%)	111 (18.3%)	142 (13.3%)	
Country of birth				0.0529
Australia	374 (82.2%)	528 (86.7%)	902 (84.8%)	
Outside of Australia	81 (17.8%)	81 (13.3%)	162 (15.2%)	
Education level (<i>N</i> = 1026)				<0.001
≤9 years	107 (24.7%)	209 (35.5%)	316 (30.9%)	
10–11 years	145 (33.4%)	218 (37.0%)	363 (35.5%)	
≥12 years	182 (41.9%)	162 (27.5%)	344 (33.6%)	
Currently employed				0.00403
Unemployed	371 (81.4%)	537 (87.9%)	908 (85.1%)	
Employed or full-time student	85 (18.6%)	74 (12.1%)	159 (14.9%)	
Unstable accommodation				<0.001
Unstable	232 (50.9%)	240 (39.3%)	472 (44.3%)	
Stable	224 (49.1%)	370 (60.7%)	594 (55.7%)	
Lives with spouse or partner (<i>N</i> = 1056)	72 (16.0%)	106 (17.6%)	178 (16.9%)	0.553
Ever incarcerated (<i>N</i> = 1054)	209 (46.4%)	350 (58.2%)	559 (53.2%)	<0.001
Ever assaulted (<i>N</i> = 1043)	205 (45.5%)	296 (50.3%)	501 (48.2%)	0.141
Age at first injection of drugs (<i>N</i> = 1064)	18.0 (6.00)	17.0 (4.00)	17.0 (5.00)	<0.001
Main illicit drug of choice (<i>N</i> = 1033)				0.489
Heroin	282 (64.1%)	385 (65.3%)	667 (64.8%)	
Amphetamines	81 (18.4%)	89 (15.1%)	170 (16.5%)	
Cannabis	55 (12.5%)	81 (13.7%)	136 (13.2%)	
Other	22 (5.00%)	35 (5.93%)	57 (5.53%)	
Injecting frequency in the past week (<i>N</i> = 1001)	5.00 (13.0)	5.00 (13.0)	5.00 (13.0)	0.166
AUDIT score (<i>N</i> = 1062)				0.362
Abstinent	197 (43.5%)	289 (47.7%)	486 (45.9%)	
8 or more (high-risk)	98 (21.6%)	127 (21.0%)	225 (21.2%)	
Less than 8 (low-risk)	158 (34.9%)	190 (31.4%)	348 (32.9%)	
Anxiety (GAD-7) (<i>N</i> = 588) ^c				0.199
Mild (score <10)	137 (57.3%)	179 (51.6%)	316 (53.9%)	
Moderate–Severe (score ≥10)	102 (42.7%)	168 (48.4%)	270 (46.1%)	
Depression (PHQ-9) (<i>N</i> = 588) ^c				0.275
Mild (score <10)	122 (51.5%)	160 (46.5%)	282 (48.5%)	
Moderate–Severe (score ≥10)	115 (48.5%)	184 (53.5%)	299 (51.5%)	
Currently on opioid agonist treatment (OAT)	206 (45.2%)	296 (48.4%)	502 (47.0%)	0.319

(Continues)

TABLE 1 (Continued)

	Not parent or caregiver (<i>N</i> = 456)	Parent or caregiver (<i>N</i> = 611)	Overall (<i>N</i> = 1067) ^b	<i>p</i> -value
Attended GP in the last 12 months (non-OAT) (<i>N</i> = 1069)	315 (69.2%)	410 (67.1%)	725 (68.0%)	0.503
Cared for by at least one parent (<i>N</i> = 572) ^d	92 (39.5%)	118 (34.9%)	210 (36.8%)	0.305
Ever removed as a child (<i>N</i> = 563) ^d	45 (19.7%)	90 (26.9%)	135 (24.0%)	0.0586
Member or child of stolen generation (<i>N</i> = 89) ^{d e}	8 (47.1%)	46 (63.9%)	54 (60.7%)	0.316

Abbreviation: AUDIT, Alcohol Use Disorder Identification Test.

^aNon-binary participants were added to the female gender category to reduce the risk of disclosure/re-identification given small numbers.

^b*N* = 1067 unless otherwise specified in the variable name column.

^cPatient Health Questionnaire (PHQ-9) and the Generalised Anxiety Disorder (GAD-7) scores were only available from 2015 onwards.

^dAdditional questions about the participant's care history were only available from June 2017 onwards.

^eThere were 119 Aboriginal or Torres Strait Islander participants recruited from June 2017 onwards.

TABLE 2 Care arrangements among parents and caregivers by gender.

	Male (<i>N</i> = 356)	Female or non-binary (<i>N</i> = 254)	Overall (<i>N</i> = 611)
Number of children—median (IQR) (<i>N</i> = 610)	2.00 (2.00)	2.00 (2.00)	2.00 (2.00)
At least one co-resident child (<i>N</i> = 606)	64 (18.1%)	96 (38.2%)	160 (26.4%)
At least one child with another parent or caregiver (<i>N</i> = 508)	243 (78.6%)	139 (70.2%)	383 (75.4%)
Child protection involvement (<i>N</i> = 605)	156 (44.4%)	182 (71.9%)	339 (56.0%)
Child ever removed from care (<i>N</i> = 451)	83 (35.0%)	136 (63.8%)	220 (48.8%)
No access rights for at least one child (<i>N</i> = 583)	106 (31.3%)	43 (17.7%)	149 (25.6%)

Abbreviation: IQR, interquartile range.

a child removed to OOHC were three times greater among women than men (adjusted odds ratio [aOR] 3.32, 95% confidence interval [CI] 1.24, 9.15). Aboriginal and Torres Strait Islander parents were more likely than non-Aboriginal and Torres Strait Islander parents to report the removal of at least one child to OOHC (aOR 2.27, 95% CI 1.07, 4.94). Parents who had ever been assaulted were more likely to report having their child/ren removed to OOHC (aOR 2.16, 95% CI 1.25, 3.79). For each additional child, the odds of child removal to OOHC increased by 40% (odds ratio 1.40, 95% CI 1.18, 1.67) (Table 3).

There was a significant interaction between gender and incarceration ($p \leq 0.001$), indicating that the association between incarceration and removal to OOHC differed by gender, with incarceration more strongly associated with removal to OOHC in women than in men (aOR 7.03, 95% CI 2.3, 22.7). The multivariable model predicted odds of child removal of 0.46 (95% CI 0.38, 0.55) for non-incarcerated women or non-binary parents compared to 0.88 (95% CI 0.81, 0.95) for incarcerated women or non-binary parents, while the odds of child removal were more

similar for non-incarcerated (0.30, 95% CI 0.20, 0.40) and incarcerated men (0.38, 95% CI 0.30, 0.45). Similarly, there was a greater protective association between being on current OAT ($p = 0.013$) and child removal to OOHC for women compared to men (odds ratio 0.34, 95% CI 0.11, 1.0). The multivariable model predicted odds of child removal rates of 0.75 (95% CI 0.67, 0.84) for women and non-binary parents not on OAT compared to 0.56 (95% CI 0.48, 0.64) for women and non-binary parents on OAT, while odds of child removal were similar for men not on OAT (0.35, 95% CI 0.27, 0.44) and on OAT (0.34, 95% CI 0.25, 0.43). There was no significant interaction between gender and experience of assault ($p = 0.238$) or the number of children ($p = 0.1234$), indicating that parents who experienced assault and parents with more children were more likely to have a child removed regardless of gender. There was also no significant interaction between gender and cohabitating with a spouse or partner ($p = 0.057$), age ($p = 0.123$), Aboriginal or Torres Strait Islander status ($p = 0.598$), employment ($p = 0.083$) or accommodation ($p = 0.179$), indicating that none of these covariates were significantly associated with child removal for any gender.

TABLE 3 Correlates for child removal to out-of-home care among parents who inject drugs; multivariable model includes interaction terms between incarceration and opioid agonist treatment with gender.

Characteristic	Bivariate				Multivariable		
	N	OR	95% CI	p-value	aOR	95% CI	p-value
Age at interview	451	0.99	0.97, 1.02	0.6	0.98	0.94, 1.02	0.2
Gender	450						
Male		—	—		—	—	
Female or non-binary		3.28	2.23, 4.84	<0.001	3.32	1.24, 9.15	0.018
Aboriginal and Torres Strait Islander status	448						
Not Aboriginal or Torres Strait Islander		—	—		—	—	
Aboriginal and/or Torres Strait Islander		3.28	1.93, 5.75	<0.001	2.27	1.07, 4.94	0.035
Country of birth	449						
Australia		—	—		—	—	
Outside of Australia		0.54	0.30, 0.94	0.032	1.17	0.52, 2.57	0.7
Education level	430						
≤9 years		—	—		—	—	
10–11 years		0.64	0.41, 1.01	0.055	0.72	0.40, 1.32	0.3
≥12 years		0.47	0.29, 0.75	0.002	0.58	0.29, 1.16	0.12
Currently employed	451						
Unemployed		—	—		—	—	
Employed or full-time student		0.30	0.15, 0.55	<0.001	0.49	0.19, 1.19	0.13
Unstable accommodation	451						
Unstable		—	—		—	—	
Stable		1.19	0.81, 1.75	0.4	1.19	0.68, 2.07	0.5
Lives with spouse or partner	446						
No		—	—		—	—	
Yes		2.18	1.32, 3.68	0.003	1.56	0.79, 3.12	0.2
Ever incarcerated	446						
No		—	—		—	—	
Yes		2.15	1.48, 3.15	<0.001	0.85	0.39, 1.83	0.7
Ever assaulted	439						
No		—	—		—	—	
Yes		2.63	1.79, 3.88	<0.001	2.16	1.25, 3.79	0.007
Age at first injection of drugs	448	0.97	0.92, 1.01	0.14	0.95	0.88, 1.01	0.13
Main illicit drug of choice	430						
Heroin		—	—		—	—	
Amphetamines		0.99	0.55, 1.75	>0.9	1.15	0.52, 2.54	0.7
Cannabis		1.06	0.61, 1.82	0.8	1.55	0.76, 3.17	0.2
Other		1.64	0.75, 3.72	0.2	2.07	0.72, 6.15	0.2
Injecting frequency in the past week	419	1.02	1.00, 1.03	0.050	0.99	0.97, 1.02	0.7
AUDIT score	450						
Abstinent		—	—		—	—	
8 or more (high-risk)		0.98	0.61, 1.59	>0.9	0.76	0.39, 1.47	0.4
Less than 8 (low-risk)		1.27	0.83, 1.95	0.3	1.28	0.70, 2.32	0.4

(Continues)

TABLE 3 (Continued)

Characteristic	Bivariate				Multivariable		
	N	OR	95% CI	p-value	aOR	95% CI	p-value
Currently on OAT	451						
No		—	—		—	—	
Yes		0.76	0.53, 1.10	0.15	1.62	0.76, 3.52	0.2
Attended GP in the last 12 months (non-OAT)	451						
No		—	—		—	—	
Yes		1.12	0.75, 1.68	0.6	0.70	0.39, 1.24	0.2
Number of children	450	1.54	1.35, 1.79	<0.001	1.40	1.18, 1.67	<0.001
Gender (female or non-binary)*Ever incarcerated					7.03	2.30, 22.7	<0.001
Gender (female or non-binary)*Currently on OAT					0.34	0.11, 1.0	0.051

Abbreviations: aOR, adjusted odds ratio; AUDIT, Alcohol Use Disorder Identification Test; CI, confidence interval; GP, general practitioner; OAT, opioid agonist treatment; OR, odds ratio.

3.2 | Intergenerational experiences of OOHC

Parents who themselves were removed to OOHC as a child were more likely to report child protection involvement ($n = 58$, 65%) compared to parents who were not removed as a child ($n = 123$, 51%). This group was also more likely to have a child removed from their care ($n = 49$, 85%) compared to parents who were not removed as a child ($n = 70$, 57%).

The sub-analysis examining the parental history of OOHC as a correlate included 156 parents of 495 children. The likelihood that their own child was removed to out-of-home care was four times greater among parents who were themselves removed as a child (odds ratio 4.05 95% CI 1.21, 15.1) (Table S4).

4 | DISCUSSION

In our cohort of people who inject drugs, those who identified as being parents were more socially disadvantaged than those who did not. Parents who inject drugs were more likely to have lower levels of educational attainment. Although the proportion of parents living in unstable housing was lower than non-parents, the proportion of parents who inject drugs living in unstable accommodation overall was considerably higher than in the general population [34]. Previous studies conducted among pregnant women in treatment for opioid use disorder in the United States and clients of a methadone maintenance program in Canada also described marked social disadvantage [19, 27]. Parents were equally as likely to experience moderate–severe levels of anxiety (48%) and depression (53%) compared to non-parents (43% and 49%

moderate–severe anxiety and depression, respectively), at rates much higher than the general population. The similar results among parents and non-parents contrast with previous research from the United States indicating higher psychiatric and psychosocial functioning among parents in treatment for opioid use disorder compared to non-parents, specifically in terms of anxiety, social connectedness, loneliness and life satisfaction [35].

We also identified high rates of child protection involvement (56%) and placement into OOHC (49%) among parents who inject drugs, with even higher proportions among women or non-binary parents, most of whom reported child protection involvement (72%) and removal of a child under their care (64%). A previous study of mothers in treatment for heroin use in Australia reported that 64% had child protection involvement and 33% had at least one child in OOHC [36]. Among mothers in substance use treatment in the United States, 47% reported child protection involvement while 34% reported having at least one child in OOHC [37]. A more recent study from Australia among parents who smoke methamphetamine reported similarly low levels of co-resident children (12% of fathers and 43% of mothers had at least one co-resident child) [22]. While most of the literature on parental drug use and child protection focuses on mothers, we identified fathers as more likely to report no access rights to their child(ren) (31.3%) compared to mothers or non-binary parents (17.7%), suggesting that removal of parental rights may be an important, but neglected issue for men who use drugs.

Child protection involvement may be appropriate for some families, given evidence suggests a small, but statistically significant detriment to child well-being associated with parental substance use [38]. While our analyses did not investigate the indications nor appropriateness of

child protection involvement or child removal among participants, the results need to be interpreted within a context of pervasive stigmatisation of people who inject drugs [21, 39]. This stigmatisation has the potential to contribute to high rates of child removal through multiple mechanisms, including increased surveillance or stricter conditions at an interpersonal level, and the use of punitive rather than supportive policies at an institutional level [17, 40].

Being female or non-binary, Aboriginal or Torres Strait Islander, experiencing assault and having a higher number of children was associated with greater odds of having a child removed into out-of-home care. Our results are consistent with previous studies in which maternal substance use features more heavily in child protection involvement than paternal substance use [41]. Further explorations of interactions by gender indicated that incarceration was associated with a higher risk of child placement into OOHC for women compared to men, while treatment on OAT was more protective for women than for men. These differing associations by gender may reflect the higher likelihood for mothers to be a child's main carer; however, they may also relate to increased scrutiny faced by women based on social and political norms and expectations about parental roles [41, 42]. Surveys of community attitudes indicate high levels of stigma towards mothers who use illicit drugs, with one-third of participants in a US survey reporting a belief that a mother who is receiving treatment for their opioid use disorder during pregnancy should be arrested, while two-thirds felt that the infant should be removed from their care [35, 43]. While our study did not specifically consider intimate partner or family violence, the high prevalence of lifetime assault and the association with child placement into OOHC may reflect this risk. There is a known association between family violence and poor child health outcomes, which may be mediated directly through the perpetration of violence against the child as well as indirectly through impacts of marital conflict [44].

Our results indicate a higher risk of child placement to OOHC among Aboriginal and Torres Strait Islander parents, consistent with longstanding disparities in patterns of child protection involvement between Aboriginal and non-Aboriginal children. This occurs in the context of a history of colonisation and forced removals with a legacy of intergenerational trauma and ongoing experiences of racism and discrimination experienced by Aboriginal and Torres Strait Islander families [45]. The 2020 Family Matters report, coordinated by a group of eminent Aboriginal and Torres Strait Islander leaders and supported by a Strategic Alliance of over 150 Aboriginal and Torres Strait Islander and non-Indigenous organisations,

proposed a roadmap to eliminate the overrepresentation of Aboriginal and Torres Strait Islander children in OOHC, which focuses on access to quality, culturally safe services and the right to self-determination [46].

A sub-analysis identified intergenerational risks for child placement to OOHC, with placement being two times higher among parents who were themselves placed in OOHC. Our findings are consistent with previous research demonstrating intergenerational continuity in responses to child removal/placement in OOHC [47, 48]. These findings are particularly relevant, given that self-reported placement in OOHC as a child was prevalent (24%) among our parents who inject drugs.

Overall, results reinforce the need for targeted psychosocial support and systems-level change addressing social disadvantage—housing, employment, family violence and mental health—to achieve equitable health outcomes for parents who inject drugs and their children [49]. Existing reviews support the role of integrated treatment programs, addressing both parenting and substance use, on parenting outcomes and to support parents to reduce their substance use if that is their goal [50–53]. Coordinated service access also impacts the health and well-being of parents and their children. Indeed, previous research indicates mothers who were treated in programs providing a high level of family-related or education/employment services were more likely to reunify with their children [54].

The gender-specific correlates highlighted in our research support previous calls for a focus on gender-focused research and gender-responsive care [55]. Given the gendered association between incarceration and child removal, progress towards drug decriminalisation is critical to addressing the high rates of child removal among women and non-binary parents identified in our cohort. Importantly, the disproportionate risk of child removal to OOHC among Aboriginal and Torres Strait Islander participants, despite adjustment for other correlates, highlights the need for tailored preventative approaches led by Aboriginal and Torres Strait Islander communities, which may include strengthening of protective cultural factors and addressing intergenerational trauma caused by government policies [56]. Approaches to addressing the needs of parents who inject drugs should also be informed by complementary existing literature identifying strengths or protective attributes within the family or environment that are associated with positive child health outcomes [16].

A key strength of this study is the inclusion of a wide range of psychosocial factors in the analysis, including parents' own experiences of OOHC. Furthermore, our study was unique in reporting on a community-based sample, which is likely to be more representative of

people who inject drugs in the community compared to previous studies recruiting from treatment programs.

However, our study has several limitations. Firstly, given the cross-sectional nature of our data, we cannot establish a temporal relationship between correlates and outcome. Ongoing follow-up with this cohort will provide further opportunity to confirm temporality. Our main outcome of removal to OOHC is based on parental report which is subject to reporting bias and may or may not align with the current Australian definition of OOHC—that is overnight care for children under the age of 18 years who are deemed unable to live with their family because of safety concerns. Similarly, we were not prescriptive with our definition of ‘caregiver’ so there may have been variation in how this was interpreted by participants. With a larger sample size, we could have considered investigating correlates for more specific outcome definitions, such as placement type (e.g., kinship or residential care) or placement at differing ages, given the differences in health and behavioural outcomes associated with these factors [57]. Our analysis was also limited by missing data, particularly for our primary outcome of child placement to OOHC (26% missing), resulting in reduced statistical power due to reduced sample size. While we are unaware of any systematic reason for the missing child placement status, there is also the risk that bias has been introduced. Finally, our dataset focusses on the characteristics of the parents who use drugs. Hence, we have limited information on the children themselves, including their ages, age at removal to OOHC or any health or behavioural outcomes.

5 | CONCLUSIONS

Our study highlights high rates of child placement in OOHC among parents who inject drugs. Sub-populations at higher risk of child removal within this cohort include women, Aboriginal and Torres Strait Islander parents, parents who have experienced assault, parents of multiple children and parents who were themselves removed to OOHC as children. Targeted services and structural changes are needed to address the systemic health and social inequities that underpin these associations. Further research is needed to establish evidence-based, culturally safe, trauma-informed, harm reduction-oriented and participant-driven policies and programs to address the high rates of out-of-home care among children of parents who inject drugs and to better support them to care for their children.

AUTHOR CONTRIBUTIONS

Jocelyn Chan undertook statistical analysis and drafted the initial manuscript, supervised by Paul Dietze. All

authors contributed to the writing of the manuscript and approved the final version.

ACKNOWLEDGEMENTS

We would like to acknowledge the contribution of the SuperMIX participants, the Burnet Institute fieldwork team and supporting community services and organisations. Open access publishing facilitated by Curtin University, as part of the Wiley - Curtin University agreement via the Council of Australian University Librarians.

FUNDING INFORMATION


Mark Stoové has received investigator-initiated research funding from Gilead Sciences and AbbVie and consultant fees from Gilead Sciences for activities unrelated to this work. Lisa Maher is funded by a National Health and Medical Research Council (NHMRC) Senior Research Fellowships. Baseline data collection for SuperMIX was funded by the Colonial Foundation Trust and the NHMRC (#545891 and #1126090), with ongoing data collection funded by the NHMRC alone. The authors gratefully acknowledge the support of the Victorian Operational Infrastructure Fund. The funders had no input into the work.

CONFLICT OF INTEREST STATEMENT

The authors declare no conflicts of interest.

ORCID

Jocelyn Chan  <https://orcid.org/0000-0002-8536-2048>

Bernadette Ward  <https://orcid.org/0000-0002-8166-6386>

Paul Dietze  <https://orcid.org/0000-0001-7871-6234>

REFERENCES

1. Degenhardt L, Peacock A, Colledge S, Leung J, Grebely J, Vickerman P, et al. Global prevalence of injecting drug use and sociodemographic characteristics and prevalence of HIV, HBV, and HCV in people who inject drugs: a multistage systematic review. *Lancet Global Health*. 2017;5:e1192–207.
2. Degenhardt L, Charlson F, Ferrari A, Santomauro D, Erskine H, Mantilla-Herrera A, et al. The global burden of disease attributable to alcohol and drug use in 195 countries and territories, 1990–2016: a systematic analysis for the global burden of disease study 2016. *Lancet Psychiatry*. 2018;5:987–1012.
3. Doab A, Fowler C, Dawson A. Factors that influence mother–child reunification for mothers with a history of substance use: a systematic review of the evidence to inform policy and practice in Australia. *Int J Drug Policy*. 2015;26:820–31.
4. Fernandez E, Lee J-S. Returning children in care to their families: factors associated with the speed of reunification. *Child Indic Res*. 2011;4:749–65.
5. Taplin S, Mattick RP. The nature and extent of child protection involvement among heroin-using mothers in treatment: high rates of reports, removals at birth and children in care. *Drug Alcohol Rev*. 2015;34:31–7.

6. Australian Institute of Health and Welfare (AIHW). Child protection Australia 2018–19. Child Welfare Series No. 72. Cat. No. CWS 74. Canberra, AIHW. 2020.
7. Australian Institute of Health and Welfare (AIHW). Child protection Australia 2020–2021. Canberra, Australia: Australian Government; 2022.
8. Gypen L, Vanderfaeillie J, de Maeyer S, Belenger L, Van Holen F. Outcomes of children who grew up in foster care: systematic-review. *Child Youth Serv Rev*. 2017;76:74–83.
9. Kääriälä A, Hiilamo H. Children in out-of-home care as young adults: a systematic review of outcomes in the Nordic countries. *Child Youth Serv Rev*. 2017;79:107–14.
10. Straatmann VS, Jackisch J, Brännström L, Almquist YB. Inter-generational transmission of out-of-home care and the role of mental health problems: findings from Stockholm birth cohort multigenerational study. *Soc Sci Med*. 2021;284:114223.
11. Tracy M, Salo M, Appleton AA. The mitigating effects of maternal social support and paternal involvement on the inter-generational transmission of violence. *Child Abuse Negl*. 2018;78:46–59.
12. Brännström L, Vinnerljung B, Almquist YB. Intergenerational transmission of placement in out-of-home care: mediation and interaction by educational attainment. *Child Abuse Negl*. 2022;123:105436.
13. Thumath M, Humphreys D, Barlow J, Duff P, Braschel M, Bingham B, et al. Overdose among mothers: the association between child removal and unintentional drug overdose in a longitudinal cohort of marginalised women in Canada. *Int J Drug Policy*. 2021;91:102977.
14. Darlington CK, Clark R, Jacoby SF, Terplan M, Alexander K, Compton P. Outcomes and experiences after child custody loss among mothers who use drugs: a mixed studies systematic review. *Drug Alcohol Depend*. 2023;251:110944.
15. Klee H. Drug-using parents: analysing the stereotypes. *Int J Drug Policy*. 1998;9:437–48.
16. Richter KP, Bammer G. A hierarchy of strategies heroin-using mothers employ to reduce harm to their children. *J Subst Abuse Treat*. 2000;19:403–13.
17. Olsen A. Punishing parents: child removal in the context of drug use. *Drug Alcohol Rev*. 2015;34:27–30.
18. Garnezy N. Stress-resistant children: the search for protective factors. Oxford: Pergamon Press; 1985.
19. Rose-Jacobs R, Trevino-Talbot M, Vibbert M, Lloyd-Travaglini C, Cabral HJ. Pregnant women in treatment for opioid use disorder: material hardships and psychosocial factors. *Addict Behav*. 2019;98:106030.
20. Heberle AE, Obus EA, Gray SAO. An intersectional perspective on the intergenerational transmission of trauma and state-perpetrated violence. *J Soc Issues*. 2020;76:814–34.
21. Weber A, Miskle B, Lynch A, Arndt S, Acion L. Substance use in pregnancy: identifying stigma and improving care. *Subst Abuse Rehabil*. 2021;12:105–21.
22. Ward B, Kippen R, Reupert A, Maybery D, Agius PA, Quinn B, et al. Parent and child co-resident status among an Australian community-based sample of methamphetamine smokers. *Drug Alcohol Rev*. 2021;40:1275–80.
23. Russell L, Gajwani R, Turner F, Minnis H. Gender, addiction, and removal of children into care. *Front Psych*. 2022;13:887660.
24. Stringer KL, Baker EH. Stigma as a barrier to substance abuse treatment among those with unmet need: an analysis of parenthood and marital status. *J Fam Issues*. 2018;39:3–27.
25. Work EC, Muftu S, Macmillan KDL, Gray JR, Bell N, Terplan M, et al. Prescribed and penalized: the detrimental impact of mandated reporting for prenatal utilization of medication for opioid use disorder. *Matern Child Health J*. 2023;27:104–12.
26. Canfield M, Radcliffe P, Marlow S, Boreham M, Gilchrist G. Maternal substance use and child protection: a rapid evidence assessment of factors associated with loss of child care. *Child Abuse Negl*. 2017;70:11–27.
27. Luo C, Sanger N, Zielinski L, Bhatt M, Shahid H, Shams I, et al. Sociodemographic characteristics of patients with children in a methadone maintenance program: a cross-sectional study. *Harm Reduct J*. 2019;16:13.
28. van den Boom W, Quiroga M, O'Keefe D, Kumar D, Hill P, Scott N, et al. Cohort profile: the Melbourne injecting drug user cohort study (SuperMIX). *Int J Epidemiol*. 2022;51:e123–30.
29. Kroenke K, Wu J, Yu Z, Bair MJ, Kean J, Stump T, et al. Patient health questionnaire anxiety and depression scale: initial validation in three clinical trials. *Psychosom Med*. 2016;78:716–27.
30. The Australian Institute of Aboriginal and Torres Strait Islander Studies. The stolen generations. 2023 Retrieved 27 February 2023. Available from: <https://aiatsis.gov.au/explore/stolen-generations>
31. Zeileis A, Hothorn T. Diagnostic checking in regression relationships. *R News*. 2002;2(3):7–10.
32. R Core Team. R: a language and environment for statistical computing. Vienna, Austria: R Foundation for Statistical Computing; 2022.
33. Arel-Bundock V. margineffects: Predictions, Comparisons, Slopes, Marginal Means, and Hypothesis Tests. R package version 0.10.0. 2023.
34. Australian Bureau of Statistics. Census of population and housing: estimating homelessness. Retrieved 19 December 2022. 2016. Available from: <https://www.abs.gov.au/statistics/people/housing/census-population-and-housing-estimating-homelessness/latest-release>
35. Sattler AF, Hooker SA, Levy R, Sherman MD. Psychosocial needs of parents engaged in treatment for opioid use disorder. *Subst Use Misuse*. 2021;56:2202–13.
36. Taplin S, Mattick RP. Mothers in methadone treatment and their involvement with the child protection system: a replication and extension study. *Child Abuse Negl*. 2013;37:500–10.
37. Grella CE, Hser Y-I, Huang Y-C. Mothers in substance abuse treatment: differences in characteristics based on involvement with child welfare services. *Child Abuse Negl*. 2006;30:55–73.
38. Kuppens S, Moore SC, Gross V, Lowthian E, Siddaway AP. The enduring effects of parental alcohol, tobacco, and drug use on child well-being: a multilevel meta-analysis. *Dev Psychopathol*. 2020;32:765–78.
39. Lipsett M, Wyant-Stein K, Mendes S, Berger E, Berkman ET, Terplan M, et al. Addressing stigma within the dissemination of research products to improve quality of care for pregnant and parenting people affected by substance use disorder. *Front Psych*. 2023;14:1199661.
40. Wolfson L, Schmidt RA, Stinson J, Poole N. Examining barriers to harm reduction and child welfare services for pregnant women and mothers who use substances using a stigma action framework. *Health Soc Care Community*. 2021;29:589–601.
41. Flacks S. Substance misuse and parenting: making drugs and gender in the family court. *Int J Law Context*. 2019;15:424–41.

42. Roy J. Children living with parental substance misuse: a cross-sectional profile of children and families referred to children's social care. *Child Fam Soc Work*. 2021;26:122–31.
43. Schiff DM, Stoltman JJK, Nielsen TC, Myers S, Nolan M, Terplan M, et al. Assessing stigma towards substance use in pregnancy: a randomized study testing the impact of stigmatizing language and type of opioid use on attitudes toward mothers with opioid use disorder. *J Addict Med*. 2022;16:77–83.
44. Chiesa AE, Kallechey L, Harlaar N, Rashaan Ford C, Garrido EF, Betts WR, et al. Intimate partner violence victimization and parenting: a systematic review. *Child Abuse Negl*. 2018;80:285–300.
45. O'Donnell M, Taplin S, Marriott R, Lima F, Stanley FJ. Infant removals: the need to address the over-representation of Aboriginal infants and community concerns of another 'stolen generation'. *Child Abuse Negl*. 2019;90:88–98.
46. SNAICC—National Voice for our children, the family matters campaign, Monash University and the University of Technology Sydney (2022). The Family Matters Report 2022.
47. Appleyard K, Berlin LJ, Rosanbalm KD, Dodge KA. Preventing early child maltreatment: implications from a longitudinal study of maternal abuse history, substance use problems, and offspring victimization. *Prev Sci*. 2011;12:139–49.
48. Berlin LJ, Appleyard K, Dodge KA. Intergenerational continuity in child maltreatment: mediating mechanisms and implications for prevention. *Child Dev*. 2011;82:162–76.
49. Carey G, Crammond B. Systems change for the social determinants of health. *BMC Public Health*. 2015;15:662.
50. Neger EN, Prinz RJ. Interventions to address parenting and parental substance abuse: conceptual and methodological considerations. *Clin Psychol Rev*. 2015;39:71–82.
51. Moreland AD, McRae-Clark A. Parenting outcomes of parenting interventions in integrated substance-use treatment programs: a systematic review. *J Subst Abuse Treat*. 2018;89:52–9.
52. Peisch V, Sullivan AD, Breslend NL, Benoit R, Sigmon SC, Forehand GL, et al. Parental opioid abuse: a review of child outcomes, parenting, and parenting interventions. *J Child Fam Stud*. 2018;27:2082–99.
53. McGovern R, Newham JJ, Addison MT, Hickman M, Kaner EFS. Effectiveness of psychosocial interventions for reducing parental substance misuse. *Cochrane Database Syst Rev*. 2021;3:CD012823.
54. Grella CE, Needell B, Shi Y, Hser YI. Do drug treatment services predict reunification outcomes of mothers and their children in child welfare? *J Subst Abuse Treat*. 2009;36:278–93.
55. Meyer JP, Isaacs KK, El-Shahawy O, Burlew AK, Wechsberg W. Research on women with substance use disorders: reviewing progress and developing a research and implementation roadmap. *Drug Alcohol Depend*. 2019;197:158–63.
56. Snijder M, Lees B, Stearne A, Ward J, Garlick Bock S, Newton N, et al. An ecological model of drug and alcohol use and related harms among Aboriginal and Torres Strait Islander Australians: a systematic review of the literature. *Prev Med Rep*. 2021;21:101277.
57. Washington T, Wrenn A, Kaye H, Priester MA, Colombo G, Carter K, et al. Psychosocial factors and behavioral health outcomes among children in Foster and Kinship care: a systematic review. *Child Youth Serv Rev*. 2018;90:118–33.

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Chan J, Ward B, Maher L, Crawford S, Stoové M, Dietze P. Parents who inject drugs: Demographics, care arrangements and correlates for child placement in out-of-home care. *Drug Alcohol Rev*. 2023. <https://doi.org/10.1111/dar.13798>